arXiv:1212.3047v1 [math.SP] 13 Dec 2012

EXTENSION OF POSITIVE DEFINITE FUNCTIONS

PALLE E.T. JORGENSEN AND ROBERT NIEDZIALOMSKI

Abstract. Let $\Omega \subset \mathbb{R}^n$ be an open subset of $\mathbb{R}^n$. We say that a function $F\colon \Omega - \Omega \to \mathbb{C}$ is positive definite if for any $x_1, \dots, x_m \in \Omega$ and any $c_1, \dots, c_m \in \mathbb{C}$ the following holds

$$\sum_{j,k=1}^{m} F(x_j - x_k) c_j \overline{c_k} \geq 0.$$

Let $F\colon \Omega - \Omega \to \mathbb{C}$ be a continuous positive definite function. We give necessary and sufficient conditions for F to have an extension to a continuous and positive definite function defined on the entire Euclidean space $\mathbb{R}^n$. The conditions are formulated in terms of strong commutativity of some certain selfadjoint operators defined on a Hilbert space associated to our positive definite function.

1. Introduction

Let $\Omega \subset \mathbb{R}^n$ be an open set and let $F\colon \Omega - \Omega \to \mathbb{C}$ be a function, where $\Omega - \Omega = \{x - y\colon x, y \in \Omega\}$. We say that F is positive definite if for any $x_1, \dots, x_m \in \Omega$ and any $c_1, \dots, c_m \in \mathbb{C}$

$$\sum_{j,k=1}^{m} F(x_j - x_k) c_j \overline{c_k} \geq 0.$$

In particular, if F is a positive definite function, then $F(0) \geq 0$ and for any $z \in \Omega - \Omega$

$$F(-z) = \overline{F(z)} \quad \text{and} \quad |F(z)| \leq F(0)$$

We consider the extension problem for continuous positive definite function defined on an open subset of the Euclidean space $\mathbb{R}^n$. By extension we always mean an extension to a continuous and positive definite function defined on the entire space $\mathbb{R}^n$. We give necessary and sufficient conditions for F to have an extension. The conditions are formulated in terms of strong commutativity of some certain selfadjoint operators defined on a Hilbert space associated to our positive definite function.

While there is a substantial literature on positive definite functions and their role in applications, the problem of extensions is less studied. Here we address this question in the case of extension from certain open subsets Ω in $\mathbb{R}^n$. For $n = 1$, this was addressed first by Krein [12], but if $n > 1$, it is known, that extensions may fail to exist, see e.g., [19] and the discussion below. On the other hand, the case of several variables is of interest in physics and in the theory of boundary values. In these areas, the distinction between local (in some open subset in $\mathbb{R}^n$) vs global (all of $\mathbb{R}^n$) concerns particles at infinity, as well as the presence of commuting observables, if we adopt the axiom of quantum theory with observables represented by selfadjoint operators in Hilbert space.

In this paper we show that for a fixed locally defined positive definite continuous function F there is an associated Hilbert space, and a system of commuting Hermitian operators with common domain, such that the existence of strongly commuting selfadjoint extensions is equivalent to extendability of F to a globally defined continuous positive definite function on $\mathbb{R}^n$.

For background references on positive definite functions and their applications, see [6], [13], [4], [3], [16], [15], [17]. In a special case, in [6], the idea of connecting spectral theory to the study of commuting selfadjoint extensions was suggested there.

Let $\Omega \subset \mathbb{R}^n$ be a symmetric ($x \in \Omega \Rightarrow -x \in \Omega\}$) and convex set. Then $\Omega - \Omega = 2\Omega = \{2x\colon x \in \Omega\}$. Krein ([12]) studies this extension problem in the case when Ω is a bounded interval $(-r, r)$, $r > 0$. He proves

---

2010 *Mathematics Subject Classification.* 47B32, 42A82, 47B25, 22D10.

Key words and phrases. Positive definite functions, reproducing-kernel Hilbert spaces, symmetric(Hermitian) operators (unbounded), commutativity, commuting selfadjoint extensions, spectral resolution, unitary representations.

that a positive definite extension always exists, but is not necessarily unique. Later Devinatz ([5]) gives necessary conditions for existence of extensions when the domain is a rectangle $(-r_1, r_1) \times (-r_2, r_2) \subset \mathbb{R}^2$. In [18] Rudin extends the result of M. Krein and proves the following. Every radial positive definite function defined on a ball $B(0, r) = \{x \in \mathbb{R}^n \colon |x| < r\}$ extends to a radial positive definite function defined on the entire space $\mathbb{R}^n$. Nussbaum ([14]) gives a characterization (similar to the theorem of Bochner, see below) of continuous positive definite functions obtaining the result of Rudin as a corollary. Moreover, in [19] the author gives an example of a bounded continuous positive definite function defined on the unit cube $I^n = \{x = (x_1, \ldots, x_n) \in \mathbb{R}^n \colon |x_i| < 1, i = 1, \ldots, n\}$ that cannot be extended. Therefore in order to be guaranteed the existence of an extension we have to impose some symmetry conditions on the function or on the domain of the function.

The problem of extending positive definite function doesn't have to be restricted to the case of a subset of the Euclidean space. The notion of a positive definite function can be introduced for example for groups and semi-groups, and the extension problem can be considered there. One article in this direction that we would like to mention, that we borrowed some ideas from, is [9], where the author studies positive definite functions defined on a Lie group and he gives necessary and sufficient conditions for existence of extensions. Moreover, the range of a positive definite function doesn't have to be the set of complex numbers. For example matrix-valued or operator-valued positive definite function have been studied.

1.1. Statement of the problem.

We have the following well-known characterization of continuous positive definite functions. Let $\Omega \subset \mathbb{R}^n$ be an open set.

Theorem 1.1. *Let $F \colon \Omega - \Omega \to \mathbb{C}$ be a continuous function. Then F is positive definite if and only if for any $\varphi \in C_0(\Omega)$ (see remark below) the following holds*

$$\int_\Omega \int_\Omega F(y - x)\varphi(x)\overline{\varphi(y)}dxdy \geq 0.$$

Remark 1.2. The theorem holds true if we replace the class $C_0(\Omega)$ of continuous functions with compact support with the class $C_0^\infty(\Omega)$ of smooth functions with compact support.

For $\varphi \in C_0^\infty(\Omega)$ we define a function $F_\varphi \colon \Omega \to \mathbb{C}$ by

$$F_\varphi(x) = \int_\Omega F(x - y)\varphi(y)dy$$

and for $\varphi, \psi \in C_0^\infty(\Omega)$ we put

$$\langle F_\varphi, F_\psi \rangle = \int_\Omega \int_\Omega F(y - x)\varphi(x)\overline{\psi(y)}dxdy.$$

We define

$$\mathcal{W} = \{F_\varphi \colon \varphi \in C_0^\infty(\Omega)\}$$

Then $\mathcal{W}$ is a complex vector space and $\langle \cdot, \cdot \rangle$ is a complex inner-product on $\mathcal{W}$ (see next section). We complete $(\mathcal{W}, \langle \cdot, \cdot \rangle)$ to get a complex Hilbert space $\mathcal{H}_F$. For $j = 1, \ldots, n$ we consider dense operators $-i\frac{\partial}{\partial x_j} \colon \mathcal{H}_F \to \mathcal{H}_F$ given by

$$\left(-i\frac{\partial}{\partial x_j}\right) F_\varphi = -iF_{\frac{\partial \varphi}{\partial x_j}}$$

for $F_\varphi \in \mathcal{W}$. The main theorem that we prove is the following.

Main Theorem. Let $F \colon \Omega - \Omega \to \mathbb{C}$ be a continuous positive definite function, where Ω is an open subset of $\mathbb{R}^n$. Then there exists an extension of F to a continuous positive definite function on the entire Euclidean space $\mathbb{R}^n$ if and only if there exist selfadjoint strongly commuting extensions of the operators $-i\frac{\partial}{\partial x_j}$.

Remark 1.3. In the above strong commuting means that the associated spectral measures commute.

Remark 1.4. The theorem is related to the following theorem of Bochner. Let $M(\mathbb{R}^n)$ be the set of all finite positive Borel measures on $\mathbb{R}^n$. For a measure $\mu \in M(\mathbb{R}^n)$ we define its Fourier transform $\widehat{\mu} \colon \mathbb{R}^n \to \mathbb{C}$ to be the following function

$$\widehat{\mu}(t) = \int_{\mathbb{R}^n} e^{i\langle t, x \rangle} d\mu(x)$$

where $\langle \cdot, \cdot \rangle$ denotes the standard inner product on $\mathbb{R}^n$. Then the following holds

Theorem 1.5. *(Bochner) Let $G\colon \mathbb{R}^n \to \mathbb{C}$ be a continuous function. Then G is a positive definite function if and only if there exists a measure $\mu \in M(\mathbb{R}^n)$ such that $G = \hat{\mu}$.*

For a proof of the Bochner theorem we refer the reader to [11].

Remark 1.6. Here we recall the definition of a positive definite tempered distribution. It can be view as a generalization of the theorem of Bochner (see [1]). Let $T \in S'$ be a tempered distribution. We say that T is positive definite if there exists a Borel measure ν on $\mathbb{R}^n$ satisfying

$$\int_{\mathbb{R}^n} \frac{d\nu(x)}{(1+|x|^2)^p} < \infty$$

for some $p \in \mathbb{N}$ such that for any test function $\varphi \in S$

$$T(\varphi) = \int_{\mathbb{R}^n} \widehat{\varphi}(x) d\nu(x).$$

1.2. **Organization of the paper.** In the next section we study the Hilbert space $\mathcal{H}_F$. We prove that the form $\langle \cdot, \cdot \rangle$ defined above is a complex inner-product on the space $\mathcal{W}$. We show existence of elements $\gamma_a \in \mathcal{H}_F$, $a \in \Omega$, that behave like Dirac delta functions. In particular, for $a, b \in \Omega$

$$F(a-b) = \int_\Omega \int_\Omega F(y-x)\gamma_b(x)\gamma_a(y)dxdy = \langle \gamma_b, \gamma_a \rangle$$

This identity plays a crucial role in the proof of the main result. Then we follow with a section where we gather some facts about orthogonal projection-valued measures. In the section 4 we show that the operators $-i\frac{\partial}{\partial x_j}$ are Hermitian. If A_j is a selfadjoint extension of the operator $-i\frac{\partial}{\partial x_j}$, we study the strongly continuous one-parameter group $U_j(t)$ of unitary transformations associated to the operator A_j, $j = 1, \ldots, n$. In section 5 we give a proof of the main theorem and in the last section we show that a positive definite function defined on an open subset of the real line always admits an extension. We show that the Hilbert space $\mathcal{H}_F$ and the reproducing kernel Hilbert space associated to F are unitary isomorphic, and we briefly talk about the connection of the extension problem with the Fuglede conjecture.

2. The Hilbert space $\mathcal{H}_F$

Let $\Omega \subset \mathbb{R}^n$ be an open set and let $F\colon \Omega - \Omega \to \mathbb{C}$ be a continuous positive definite function. Recall that for $\varphi \in C_0^\infty(\Omega)$ the function $F_\varphi\colon \Omega \to \mathbb{C}$ is given by

$$F_\varphi(x) = \int_\Omega F(x-y)\varphi(y)dy.$$

and for $\varphi, \psi \in C_0^\infty(\Omega)$ we define

$$\langle F_\varphi, F_\psi \rangle = \int_\Omega \int_\Omega F(y-x)\varphi(x)\overline{\psi(y)}dxdy$$

Denote by F^e the extension of F to $\mathbb{R}^n$ by putting $F^e = 0$ in the complement of $\Omega - \Omega$. Then F_φ is the convolution of F^e with φ i.e. for $x \in \Omega$

$$F_\varphi(x) = F^e \star \varphi(x) = \int_{\mathbb{R}^n} F^e(x-y)\varphi(y)dy$$

and for $\varphi, \psi \in C_0^\infty(\Omega)$

$$\langle F_\varphi, F_\psi \rangle = \int_{\mathbb{R}^n} F^e \star \varphi(y)\overline{\psi(y)}dx = \int_{\mathbb{R}^n} F_\varphi(x)\overline{\psi(x)}dx \tag{1}$$

We note that for any $\varphi \in C_0^\infty(\Omega)$ the function $F^e \star \varphi$ is smooth and it has compact support when Ω is a bounded set.

Lemma 2.1. *The form $\langle \cdot, \cdot \rangle$ is a complex inner-product on the space $\mathcal{W}$.*

Proof. We need to show that $\langle \cdot, \cdot \rangle$ is positive definite i.e.

$$\langle F_\varphi, F_\varphi \rangle = 0 \Rightarrow F_\varphi = 0$$

We denote by (ϕ_ϵ) a smoothing kernel i.e we take a smooth non-negative function ϕ with support contained in the unit ball $B_1(0) = \{x \in \mathbb{R}^n : |x| < 1\}$ satisfying $\int \phi = 1$, and we define $\phi_\epsilon(x) = 1/\epsilon^n \phi(x/\epsilon)$. Moreover, for any $a \in \mathbb{R}^n$ we put $\phi_{a,\epsilon}(x) = \phi_\epsilon(x - a)$.

Let $F_\varphi \in \mathcal{W}$ be such that $\langle F_\varphi, F_\varphi \rangle = 0$. We take $\lambda \in \mathbb{C}$ and $F_\psi \in \mathcal{W}$ and we consider the quantity

$$\langle F_\varphi + \lambda F_\psi, F_\varphi + \lambda F_\psi \rangle,$$

which we know is non-negative by (1.1). Now we proceed as in the proof of the Cauchy-Schwarz inequality to get that

$$\langle F_\varphi, F_\psi \rangle = 0$$

Indeed, we compute

$$\begin{aligned} 0 &\le \langle F_\varphi + \lambda F_\psi, \varphi + \lambda F_\psi \rangle \\ &= \langle F_\varphi, F_\varphi \rangle + \overline{\lambda} \langle F_\varphi, F_\psi \rangle + \lambda \langle F_\psi, F_\varphi \rangle + |\lambda|^2 \langle F_\psi, F_\psi \rangle \\ &= 2\mathrm{Re}(\overline{\lambda} \langle F_\varphi, F_\psi \rangle) + |\lambda|^2 \langle F_\psi, F_\psi \rangle \end{aligned}$$

Taking $\Theta \in [0, 2\pi)$ such that $e^{-i\Theta} \langle F_\varphi, F_\psi \rangle \ge 0$ and putting $\lambda = te^{i\Theta}$ for $t \in \mathbb{R}$ we get that

$$0 \le 2t |\langle F_\varphi, F_\psi \rangle| + t^2 \langle F_\psi, F_\psi \rangle$$

Therefore, since $t \in \mathbb{R}$ is arbitrary, we conclude that $\langle F_\varphi, F_\psi \rangle = 0$. The fact that $F_\varphi = 0$ follows from the lemma below. □

Lemma 2.2. *Put* $F_{a,\epsilon} = F_{\phi_{a,\epsilon}}$. *For any* $a \in \Omega$

$$\langle F_\varphi, F_{a,\epsilon} \rangle \to F_\varphi(a) \tag{2}$$

Proof. We have

$$\begin{aligned} |\langle F_\varphi, F_{a,\epsilon} \rangle - F_\varphi(a)| &= \left| \int_\Omega \int_\Omega F(y-x)\varphi(x)\phi_{a,\epsilon}(y)dxdy - \int_\Omega F(a-x)\varphi(x)dx \right| \\ &= \left| \int_\Omega \int_\Omega (F(y-x) - F(a-x))\varphi(x)\phi_{a,\epsilon}(y)dxdy \right| \end{aligned}$$

Now we use continuity of the function F. □

The Hilbert space $\mathcal{H}_F$ as the completion of the space $(\mathcal{W}, \langle \cdot, \cdot \rangle)$ consists of equivalence classes of Cauchy sequences of $\mathcal{W}$, where the equivalence relation is given by

$$(F_{\varphi_n}) \sim (F_{\psi_n}) \Leftrightarrow \|F_{\varphi_n} - F_{\psi_n}\| \to 0$$

for two Cauchy sequences (F_{φ_n}) and (F_{ψ_n}). The inner-product in $\mathcal{H}_F$ is given by

$$\langle [(F_{\varphi_n})]_\sim, [(F_{\psi_n})]_\sim \rangle = \lim_{n \to \infty} \langle F_{\varphi_n}, F_{\psi_n} \rangle$$

Moreover, the space $\mathcal{W}$ is a dense subset of $\mathcal{H}_F$ under the identification of an element $F_\varphi \in \mathcal{W}$ with the constant Cauchy sequence (F_φ). The following holds.

Lemma 2.3. *For any* $a \in \Omega$ *the sequence* $(F_{a,1/n})$ *is Cauchy in* $\mathcal{W}$. *We denote* $\gamma_a = [(F_{a,1/n})]_\sim \in \mathcal{H}_F$.

Proof. The proof is similar to the proof of the lemma (2.2). □

Combining lemma 2.2 and lemma 2.3 we get that for any $F_\varphi \in \mathcal{W}$ and any $a \in \Omega$

$$F_\varphi(a) = \langle F_\varphi, \gamma_a \rangle \tag{3}$$

We call (3) the reproducing identity and we call the collection $\{\gamma_a : a \in \Omega\}$ the reproducing kernel for $\mathcal{H}_F$ (see the last section). Therefore the set span $\{\gamma_a : a \in \Omega\}$ is a dense subset of $\mathcal{H}_F$. We also get that

$$\langle \gamma_b, \gamma_a \rangle = F(a - b) \tag{4}$$

for any $a, b \in \Omega$.

3. PROJECTION-VALUED MEASURES

Let X be a metric space and let $\mathcal{B} = \mathcal{B}_X$ be the Borel σ-algebra of subsets of X. Let H be a Hilbert space and let $\mathrm{Proj}(H)$ be the set of orthogonal projections P on H i.e. $P = P^2 = P^\star$. A mapping $E\colon \mathcal{B} \to \mathrm{Proj}(H)$ is said to be an orthogonal projection-valued measure if

(i) For any $f \in H$ the function $\mu_f\colon \mathcal{B} \to [0,\infty)$ given by

$$\mu_f(\Delta) = \|E(\Delta)f\|^2 = \langle E(\Delta)f, f\rangle = \langle f, E(\Delta)f\rangle$$

is a finite Borel measure with

$$\mu_f(X) = \int_X d\mu_f = \|f\|^2$$

(ii) For any $\Delta_1, \Delta_2 \in \mathcal{B}$ we have that

$$E(\Delta_1 \cap \Delta_2) = E(\Delta_1)E(\Delta_2)$$

We will need the following.

Lemma 3.1. *Let H be a Hilbert space, let $n > 1$, and let $\{A_j\}_{j=1}^n$ be a system of selfadjoint operators (generally unbounded), each with dense domain in H, and let E_j be the orthogonal projection-valued measure associated to A_j i.e*

$$A_j = \int_{\mathbb{R}} x E_j(dx).$$

Then the following conditions are equivalent:

(i) *The operators $\{A_j\}_{j=1}^n$ strongly commute*

(ii) *For $z_j \in \mathbb{C}$ with $Im\, z_j \neq 0$, $j = 1,\ldots,n$, the system of bounded operators $\{(z_j I - A_j)^{-1}\}_{j=1}^n$ commutes*

(iii) *For $t = (t_1,\ldots,t_n) \in \mathbb{R}^n$ put*

$$U(t) = e^{it_1A_1}e^{it_2A_2}\cdot \ldots \cdot e^{it_nA_n} = \prod_{j=1}^n \int_{\mathbb{R}} e^{it_jx_j} dE_j(x_j)$$

Then $\{U(t)\}_{t\in\mathbb{R}^n}$ is a strongly continuous unitary representation of $(\mathbb{R}^n,+)$ on H.

(iv) *There is a orthogonal projection-valued measure E defined on the Borel σ-algebra $\mathcal{B}_n = \mathcal{B}_{\mathbb{R}^n}$ such that for all sets $\Delta_1 \in \mathcal{B}_1$*

$$E_j(\Delta) = E(\mathbb{R} \times \ldots \times \Delta \times \ldots \times \mathbb{R}).$$

Proof. See [8],[7],[10]. ☐

4. THE OPERATORS $-i\frac{\partial}{\partial x_j}$.

Lemma 4.1. *The operators $-i\frac{\partial}{\partial x_j}\colon \mathcal{W} \to \mathcal{W}$, $j = 1,\ldots,n$, are Hermitian i.e. for any $F_\varphi, F_\psi \in \mathcal{W}$*

$$\left\langle \left(-i\frac{\partial}{\partial x_j}\right) F_\varphi, F_\psi \right\rangle = \left\langle F_\varphi, \left(-i\frac{\partial}{\partial x_j}\right) F_\psi \right\rangle \tag{5}$$

Proof. We compute using the integration by parts formula

$$\begin{aligned}\left\langle \left(-i\frac{\partial}{\partial x_j}\right) F_\varphi, F_\psi \right\rangle &= -i\int_{\mathbb{R}^n} \left(F^e \star \frac{\partial\varphi}{\partial x_j}(x)\right)\overline{\psi(x)}dx = -i\int_{\mathbb{R}^n} \frac{\partial\varphi}{\partial x_j}(F^e \star \varphi(x))\overline{\psi(x)}dx \\ &= i\int_{\mathbb{R}^n} (F^e \star \varphi(x))\overline{\frac{\partial\psi}{\partial x_j}}(x)dx = \left\langle F_\varphi, \left(-i\frac{\partial}{\partial x_j}\right) F_\psi \right\rangle\end{aligned}$$

☐

Let $A_j\colon D_j \subset \mathcal{H}_F \to \mathcal{H}_F$ be a selfadjoint extension of the operator

$$-i\frac{\partial}{\partial x_j}\colon \mathcal{W} \subset \mathcal{H}_{\mathcal{F}} \to \mathcal{W} \subset \mathcal{H}_{\mathcal{F}}.$$

Let E_j be the orthogonal projection-valued measure associated to A_j and let U_j be the strongly continuous one-parameter group of unitary operators defined by A_j i.e

$$U_j(t) = e^{itA_j} = \int_{\mathbb{R}} e^{itx} dE_j(x)$$

for any $t \in \mathbb{R}$. Then A_j is uniquely determined by U_j and for $f \in D_j$ and $t \in \mathbb{R}$ we have that $U_j(t)f \in D_j$. Moreover

$$D_j = \{f \in \mathcal{H}_F \colon \frac{d}{dt}|_{t=0} U(t)f = \lim_{t\to 0} \frac{U(t)f - f}{t} \text{ exists}\}$$

and for $f \in D_j$ and $s \in \mathbb{R}$.

$$A_j f = -i\frac{d}{dt}|_{t=0} U_j(t)f \quad \text{and} \quad \frac{d}{dt}|_{t=s} U_j(t)f = iU_j(s)A_j f = iA_j U_j(s)f$$

We will need the following. We denote by $(e_1, \dots, e_n)$ the standard basis of $\mathbb{R}^n$.

Lemma 4.2. *Let $a, b \in \mathbb{R}$ with $a < b$ be such that $[ae_j, be_j] = \{a(1-t)e_j + bte_j \colon 0 \le t \le 1\} \subset \Omega$. Then*

$$U_j(b-a)\gamma_{ae_j} = \gamma_{be_j}. \tag{6}$$

Proof. Let $a, b \in \mathbb{R}$ be such that $[ae_j, be_j] \in \Omega$, let $F_\varphi \in \mathcal{W}$, and let (ϕ_ϵ) be a smoothing kernel. For each $\epsilon > 0$ we consider $g_\epsilon : [0, b-a] \to \mathbb{C}$ given by

$$g(t) = \langle F_\varphi, U_j(b-a-t)F_{(a+t)e_j,\epsilon}\rangle$$

Then

$$\begin{aligned}\langle F_\varphi, F_{be_j,\epsilon} - U_j(b-a)F_\epsilon\rangle &= g(b-a) - g(0) = \int_0^a g'(s)ds \\ &= \int_0^a \langle F_\varphi, \left(\frac{d}{dt}|_{t=s} U_j(b-a-t)\right) F_{(a+t)e_j,\epsilon} + U_j(b-a-s)\frac{d}{dt}|_{t=s} F_{(a+t)e_j,\epsilon}\rangle ds \\ &= \int_0^a \langle F_\varphi, U_j(b-a-s)\left(-\frac{\partial}{\partial x_j} F_{(a+t)e_j,\epsilon} + \frac{d}{dt}|_{t=s} F_{(a+t)e_j,\epsilon}\right)\rangle ds = 0,\end{aligned}$$

since

$$-\frac{\partial}{\partial x_j} F_{(a+t)e_j,\epsilon} + \frac{d}{dt}|_{t=s} F_{(a+t)te_j,\epsilon} = 0$$

We get that

$$F_{be_j,\epsilon} = U_j(b-a)F_{ae_j,\epsilon} \tag{7}$$

Taking the limit in (7) when $\epsilon \to 0$ yields (6) ☐

5. Proof of the Main Theorem

Proof of the Main Theorem. ($\Leftarrow$) Suppose that there exist self-adjoint commuting operators $A_j \colon D_j \subset \mathcal{H}_F \to \mathcal{H}_F$ that extend the operators $-i\frac{\partial}{\partial x_j} \colon \mathcal{W} \subset \mathcal{H}_{\mathcal{F}} \to \mathcal{W} \subset \mathcal{H}_{\mathcal{F}}$, $j = 1, \dots, n$. Let E_j be the orthogonal projection-valued measure associated to A_j. We define the following strongly continuous n-parameter group of unitary transformations of $\mathcal{H}_F$. For $t = (t_1, \dots, t_n) \in \mathbb{R}^n$ define

$$U(t) = e^{i\sum_{j=1}^n t_j A_j} = \prod_{j=1}^n U_j(t_j) = \prod_{j=1}^n \int_{\mathbb{R}} e^{it_j x_j} dE_j(x_j) = \int_{\mathbb{R}^n} e^{i\langle t,x\rangle} dE(x),$$

(see lemma 3.1). We fix $x_0 \in \Omega$ and we define a function $\widetilde{F} \colon \mathbb{R}^n \to \mathbb{C}$ by

$$\widetilde{F}(t) = \langle \gamma_{x_0}, U(t)\gamma_{x_0}\rangle$$

Hence, if we denote by μ the finite positive Borel measure

$$\mathcal{B}_n \ni \Delta \to \langle \gamma_{x_0}, E(\Delta)\gamma_{x_0}\rangle,$$

then

$$\widetilde{F}(t) = \int_{\mathbb{R}^n} e^{i\langle t,x\rangle} d\mu(x) = \hat{\mu}(t)$$

Therefore $\widetilde{F}$ is a continuous positive definite function. We will show that $\widetilde{F}$ is an extension of our function F. Since $U_j(t) = U(te_j)$ for $t \in \mathbb{R}$, we get by (6) for $a - b \in \Omega - \Omega$,

$$\widetilde{F}(a-b) = \langle \gamma_{x_0}, U(a-b)\gamma_{x_0}\rangle = \langle U(b)\gamma_{x_0}, U(a)\gamma_{x_0}\rangle = \langle \gamma_b, \gamma_a\rangle = F(a-b)$$

($\Rightarrow$) Let $\widetilde{F}\colon \mathbb{R}^n \to \mathbb{C}$ be a continuous positive definite function that extends our function F. Then by the theorem of Bochner there exists a finite Borel measure μ such that $F = \hat{\mu}$. For any $F_\varphi, F_\psi \in \mathcal{W}$ we have

$$\begin{aligned}\langle F_\varphi, F_\psi\rangle &= \int_{\mathbb{R}^n}\int_{\mathbb{R}^n} F(y-x)\varphi(x)\overline{\psi(y)}dxdy \\ &= \int_{\mathbb{R}^n}\left(\int_{\mathbb{R}^n} e^{-i\langle t,x\rangle}\varphi(x)dx\right)\left(\overline{\int_{\mathbb{R}^n} e^{-i\langle t,y\rangle}\psi(y)dy}\right)d\mu(t)\end{aligned}$$

We denote for $\varphi \in C_0^\infty(\mathbb{R}^n)$

$$\widehat{\varphi}(t) = \int_{\mathbb{R}^n} e^{-i\langle t,x\rangle}\varphi(x)dx$$

for $t \in \mathbb{R}^n$. Then

$$\langle F_\varphi, F_\psi\rangle = \int_{\mathbb{R}^n} \widehat{\varphi}(t)\overline{\widehat{\psi}(t)}d\mu(t) \tag{8}$$

Therefore the map

$$\mathcal{W} \ni F_\varphi \mapsto \widehat{\varphi} \in L^2(d\mu)$$

is a linear isometry, which can be extended to a linear isometry $W\colon \mathcal{H}_F \to L^2(d\mu)$. For $t \in \mathbb{R}^n$ we consider the multiplication operator

$$e^{2\pi i\langle t,\cdot\rangle}\colon L^2(d\mu) \to L^2(d\mu) \tag{9}$$

We obtain a strongly continuous n-parameter group $V(t)$ of unitary transformations of $L^2(d\mu)$ i.e.

$$V(t)f = e^{2\pi i\langle t,\cdot\rangle}f(\cdot)$$

Therefore $W(t) = W^\star V(t)W$ is a strongly continuous n-parameter group of unitary transformations on $\mathcal{H}_F$. We put

$$D_j = \{f \in \mathcal{H}_F\colon \frac{\partial}{\partial t_j}|_{t=0}W(t)f = \lim_{t\to 0}\frac{W(t)f - f}{t} \text{ exists}\}$$

and we define an operator $A_j\colon D_j \to \mathcal{H}_F$ by

$$A_j f = -i\frac{\partial}{\partial t_j}|_{t=0}W(t)f$$

Then, it is well-known, that the operators A_j are selfadjoint and that they strongly commute. Moreover they extend the operators $-i\frac{\partial}{\partial x_j}\colon \mathcal{W} \to \mathcal{W}$. Indeed, for $F_\varphi \in \mathcal{W}$ we have

$$\begin{aligned}A_jF_\varphi &= -i\frac{\partial}{\partial t_j}|_{t=0}W(t)F_\varphi = -iW^\star\left(\frac{\partial}{\partial t_j}|_{t=0}V(t)\widehat{\varphi}\right) \\ &= -iW^\star(x \to 2\pi i x_j\widehat{\varphi}(x)) = -i\frac{\partial}{\partial x_j}F_\varphi\end{aligned}$$

This finishes the proof. □

6. Remarks

We collect some corollaries to the main theorem.

Corollary 6.1. *Let $CE(F) = \{A = (A_1, \dots, A_n)\}$ be the set of all strongly commuting extensions A_j, $j = 1, \dots, n$, of the operators $-i\frac{\partial}{\partial x_j}\colon \mathcal{W} \to \mathcal{W}$, and let $M(F)$ be the set of all finite Borel measures μ on $\mathbb{R}^n$ such that $F(z) = \widehat{\mu}(z)$ for $z \in \Omega - \Omega$. Then there exists a bijective correspondence $CE(F) \leftrightarrow M(F)$.*

Corollary 6.2. *A continuous positive definite function $F\colon \Omega - \Omega \to \mathbb{C}$, with $\Omega \subset \mathbb{R}$ being an open set, has always an extension to a continuous positive definite function defined on the entire real line $\mathbb{R}$.*

Proof. We need to show that the operator

$$-i\frac{d}{dx}\colon \mathcal{W}\subset\mathcal{H}_F\to\mathcal{W}\subset\mathcal{H}_F$$

has a selfadjoint extension. Since any open subset of $\mathbb{R}$ is a countable union of disjoint open intervals, we may assume that Ω is an interval (p,r) for some $p<r$. We define $J\colon (p,r)\to(p,r)$ by $J(t)=-t+p+r$. For $\varphi\in C_0^\infty(\Omega)$ put $\varphi_J\colon\Omega\to\mathbb{C}$ by $\varphi_J(t)=\overline{\varphi(J(t))}$ and then set $JF_\varphi=F_{\varphi_J}$. Then $J\colon\mathcal{W}\to\mathcal{W}$ is a conjugation i.e. J is conjugate-linear $J(iF_\varphi)=-iJF_\varphi$, $J^2=\mathrm{Id}$, and $\|JF_\varphi\|=\|F_\varphi\|$ for any $F_\varphi\in\mathcal{W}$. Moreover $J\circ(-i\frac{\partial}{\partial x_j})=(-i\frac{\partial}{\partial x_j})\circ J$, $j=1,\dots,n$. This follows from the fact that J is conjugate-linear and that

$$\frac{\partial\varphi_J}{\partial x_j}=-\left(\frac{\partial\varphi}{\partial x_j}\right)_J$$

Thus $-i\frac{\partial}{\partial x_j}$, as a Hermitian operator that commutes with a conjugation has an extension to a selfadjoint operator, by a theorem of von Neumann. ☐

Let μ be a finite positive Borel measure on $\mathbb{R}^n$. We define a representation $\{V_\mu(t)\}_{t\in\mathbb{R}^n}$ on $L^2(d\mu)$ by

$$(V_\mu(t)f)(x)=e(x\cdot t)f(x)$$

where $f\in L^2(d\mu)$, $x,t\in\mathbb{R}^n$, and

$$e(x\cdot t)=e^{2\pi i\sum_{j=1}^n x_jt_j}$$

We have the following.

Corollary 6.3. *Let $\Omega\subset\mathbb{R}^n$ be an open set and let $F\colon\Omega-\Omega\to\mathbb{C}$ be a continuous positive definite function. Then F is extendible to a continuous positive definite function $\widetilde{F}$ defined on $\mathbb{R}^n$ if and only if there exists a unitary representation $\{U(t)\}_{t\in\mathbb{R}^n}$ on $\mathcal{H}_F$ such that $(U(t),\mathcal{H}_F)$ is unitary equivalent to the representation $(V_\mu,L^2(d\mu))$, where μ is a positive definite measure such that $\widetilde{F}=\widehat{\mu}$.*

6.1. **Connection with the Fuglede conjecture (see** [6]**).** We can extend the notion of a positive definiteness from function to distributions. The definition was introduced by L. Schwartz ([20]). Let $\Omega\subset\mathbb{R}^n$ be an open set and let $T\in\mathcal{D}'(\Omega-\Omega)$ be a distribution. For a function $\varphi\in C_0^\infty(\Omega)$ and $x\in\Omega$ we define $\varphi_x\colon x-\Omega\to\mathbb{C}$ by $\varphi_x(x-y)=\varphi(y)$, where $y\in\Omega$. Since $x-\Omega\subset\Omega-\Omega$ we can view φ_x as an element of the space $C_0^\infty(\Omega-\Omega)$. We put

$$T_\varphi(x)=T(\varphi_x)$$

for $x\in\Omega$. Then T_φ is a smooth function on Ω (compare with the convolution of a distribution with a test function). We say that the distribution T is positive definite if for any $\varphi\in C_0^\infty(\Omega)$ we have that

$$\int_\Omega T_\varphi(x)\overline{\varphi(x)}dx\geq 0$$

We consider the delta Dirac function $\delta=\delta_0$. For $\varphi\in C^\infty(\Omega)$ we have that $\delta_\varphi=\varphi_x(0)=\varphi_x(x-x)=\varphi(x)$. Therefore the delta Dirac distribution δ is positive definite. We can carry over the construction of the Hilbert space $\mathcal{H}_F$, where $F\colon\Omega-\Omega\to\mathbb{C}$ is a continuous positive definite function, to the case of the delta Dirac function δ. We get the Hilbert space $\mathcal{H}_\delta$ (see (1)). Since $\delta_\varphi=\varphi$, we see that $\mathcal{H}_\delta=L^2(\Omega)$. If $\Omega\subset\mathbb{R}$ then the dense operator $-i\frac{d}{dx}\colon L^2(\Omega)\to L^2(\Omega)$ has a selfadjoint extension. Therefore we have the associated projection valued-measure E. Denote $\Lambda=\mathrm{spec}(E)=\{\lambda\in\mathbb{R}\colon E(\{\lambda\})>0\}$, and let's look at the following two examples. (1) Let $\Omega=(0,1)\cup(2,3)\subset\mathbb{R}$. Then $\Lambda=\{0,1/4\}+\mathbb{Z}$ and the corresponding eigenspaces have dimension one. (2) Let $\Omega=(0,1)\cup(3,5)\subset\mathbb{R}$. Then $\Lambda=(1/2)\mathbb{Z}$ and the corresponding eigenspaces have dimension one or two.

6.2. **Reproducing kernel Hilbert space associated to a positive definite function.** There is a construction that associate a reproducing kernel Hilbert space to a positive definite kernel (see [2]). In our case, if $F\colon\Omega-\Omega\to\mathbb{C}$ is a positive definite function, where Ω is an open subset of $\mathbb{R}^n$, we can associate a reproducing kernel Hilbert space in the same way. Namely, for each $a\in\Omega$ we define a function $k_a\colon\Omega\to\mathbb{R}$ by

$$k_a(x)=F(x-a)$$

Then the space $W_F = \operatorname{span}\{k_a \colon a \in \Omega\}$ together with the inner-product

$$\langle \sum_{j=1}^{m} c_j k_{a_j}, \sum_{j=1}^{m} d_j k_{a_j} \rangle_F = \sum_{j,k=1}^{m} F(a_k - a_j) c_j \overline{d_k}$$

is a complex pre-Hilbert space. Its completion H_F is called the reproducing kernel Hilbert space associated to F. We note that

$$\langle k_a, k_b \rangle_F = F(b-a)$$

Since

$$\sum_{j=1}^{m} c_j \gamma_{a_j} = 0 \Leftrightarrow \sum_{j,k=1}^{m} F(a_k - a_j) c_j \overline{c_k} = 0 \Leftrightarrow \sum_{j=1}^{m} c_j k_{a_j} = 0$$

the assignment

$$\gamma_a \mapsto k_a$$

gives rise to a well-defined map between Hilbert spaces $\mathcal{H}_F$ and H_F, which is a unitary isomorphism.

We can repeat the above construction of the reproducing kernel Hilbert space in the case of the Dirac delta function i.e we identify δ with the positive definite function $\chi = \chi_0$ given by $\chi(0) = 1$ and $\chi(x) = 0$ for $x \neq 0$. Then $H_\delta = l^2(\Omega)$. Since we have that $\mathcal{H}_\delta = L^2(\Omega)$, we conclude that $\mathcal{H}_\delta \neq H_\delta$.

References

[1] Alpay, Daniel; Jorgensen, Palle; Levanony, David A class of Gaussian processes with fractional spectral measures. J. Funct. Anal. 261 (2011), no. 2, 507-541

[2] Aronszajn, N. Theory of reproducing kernels. Trans. Amer. Math. Soc. 68 (1950), 337-404

[3] Bohnstengel, Jana; Jorgensen, Palle Geometry of spectral pairs. Anal. Math. Phys. 1 (2011), no. 1, 69-99

[4] Cho, Ilwoo; Jorgensen, Palle E. T. Moment computations for graphs with fractal property. J. Appl. Math. Comput. 37 (2011), no. 1-2, 377-406

[5] Devinatz, Allen On the extensions of positive definite functions. Acta Math. 102 (1959), 109-134

[6] Fuglede, Bent Commuting self-adjoint partial differential operators and a group theoretic problem. J. Functional Analysis 16 (1974), 101-121

[7] Jorgensen, Palle E. T. Operators and representation theory. Canonical models for algebras of operators arising in quantum mechanics. North-Holland Mathematics Studies, 147. Notas de Matemtica [Mathematical Notes], 120. North-Holland Publishing Co., Amsterdam, 1988

[8] Jorgensen, Palle E. T. Extensions and index of Hermitian representations. Publ. Res. Inst. Math. Sci. 25 (1989), no. 6, 923-945

[9] Jorgensen, Palle E. T. Integral representations for locally defined positive definite functions on Lie groups. Internat. J. Math. 2 (1991), no. 3, 257-286

[10] Palle E. T. Jorgensen; Feng Tian Functional Analysis, arXiv:1003.1117

[11] Katznelson, Yitzhak An introduction to harmonic analysis. Third edition. Cambridge Mathematical Library. Cambridge University Press, Cambridge, 2004

[12] Krein, M. Sur le probleme du prolongement des fonctions hermitiennes positives et continues. C. R. (Doklady) Acad. Sci. URSS (N.S.) 26 (1940), 17-22

[13] Nelson, Edward Analytic vectors. Ann. of Math. (2) 70 (1959), 572-615

[14] Nussbaum, A. E. Integral representation of functions and distributions positive definite relative to the orthogonal group. Trans. Amer. Math. Soc. 175 (1973), 335-387

[15] Parthasarathy, K. R. Extremal quantum states in coupled systems. Ann. Inst. H. Poincar Probab. Statist. 41 (2005), no. 3, 257-268

[16] Parthasarathy, K. R. Probability measures on metric spaces. Reprint of the 1967 original. AMS Chelsea Publishing, Providence, RI, 2005

[17] Parthasarathy, K. R.; Varadhan, S. R. S. Extension of stationary stochastic processes. Teor. Verojatnost. i Primenen. 9 (1964), 72-78

[18] Rudin, Walter The extension problem for positive-definite functions. Illinois J. Math. 7 (1963), 532-539

[19] Rudin, Walter An extension theorem for positive-definite functions. Duke Math. J. 37 (1970), 49-53

[20] Schwartz, Laurent Sous-espaces hilbertiens d'espaces vectoriels topologiques et noyaux associes (noyaux reproduisants). J. Analyse Math. 13 (1964), 115-256

Department of Mathematics, The University of Iowa, 14 MacLean Hall, Iowa City, IA-52242, USA
E-mail address: `palle-jorgensen@uiowa.edu, robert-niedzialomski@uiowa.edu`